

NSW Threatened Species Scientific Committee

Notice of Preliminary Determination

The NSW TSSC has also made a Preliminary Determination to support a proposal to list the shrub *Zieria odorifera* subsp. *warrabahensis* Duretto & P.I.Forst. as a CRITICALLY ENDANGERED SPECIES in Part 1 of Schedule 1 of the Act.

A copy of these Determinations, which contains the reasons for the determinations, may be obtained free of charge on the Internet www.environment.nsw.gov.au, by contacting the NSW Threatened Species Scientific Committee, PO Box 1967 Hurstville BC 1481. Tel: (02) 9585 6940 or Fax (02) 9585 6606, or in person at the Office of Environment and Heritage Information Centre, Level 14, 59-61 Goulburn Street, Sydney. Copies of the determination may also be obtained from National Parks and Wildlife Service Area Offices and Visitor Centres, subject to availability.

How to make a submission on the Preliminary Determination

The NSW TSSC welcomes public involvement in the assessment process and places preliminary determinations on public exhibition on the NSW TSSC pages on the Office of Environment and Heritage (OEH) website. This public exhibition provides an opportunity for the public to comment on this preliminary determination as well as provide any additional information that is relevant to the assessment of the species.

Postal submissions regarding this Preliminary Determination may be sent to:

Suzanne Chate
NSW Threatened Species Scientific Committee
PO Box 1967
Hurstville BC 1481.

Email submissions in Microsoft Word or PDF formats may be sent to:
scientific.committee@environment.nsw.gov.au

Submissions close 19th October 2018.

What happens next?

After considering any submissions received during the public exhibition period the NSW TSSC will make a Final Determination and a notice will be placed on the OEH website to announce the outcome of the assessment. If the Final Determination is to list a species, then it will be added to the threatened species Schedules when the Final Determination is published on the legislation website. www.legislation.nsw.gov.au.

Privacy information

The information you provide in your submission may be used by the NSW TSSC in the assessment to determine the conservation status and listing or delisting of threatened or extinct species, threatened populations and threatened or collapsed ecological communities or to assess key threatening processes.

The NSW TSSC may be asked to share information on assessments with NSW Government agencies, the Commonwealth Government and other State and Territory governments to collaborate on national threatened species assessments using a common assessment method and to assist in the management of species and ecological communities.

NSW Threatened Species Scientific Committee

If your submission contains information relevant to the assessment it may be provided to state and territory government agencies and scientific committees as part of this collaboration.

If you wish your identity and personal information in your submission to be treated as confidential you must:

- ***request your name be treated as confidential***, and
- ***not include any of your personal information in the main text of the submission or attachments so that it can be easily removed.***

Dr Marco Duretto
Chairperson
NSW Threatened Species Scientific Committee

NSW Threatened Species Scientific Committee

Exhibition period: 24/08/18 – 19/10/18

Preliminary Determination

The NSW Threatened Species Scientific Committee, established under the *Biodiversity Conservation Act 2016* (the Act), has made a Preliminary Determination to list the shrub *Zieria odorifera* subsp. *warrabahensis* Duretto & P.I.Forst. as a CRITICALLY ENDANGERED SPECIES in Part 1 of Schedule 1 of the Act. Listing of Critically endangered species is provided for by Part 4 of the Act.

The NSW Threatened Species Scientific Committee has found that:

1. *Zieria odorifera* subsp. *warrabahensis* Duretto & P.I.Forst. (family Rutaceae) is described as an erect shrub to 40 cm tall; branches ridged, with a moderately dense indumentum of mainly stellate and some bifid hairs between the ± glabrous leaf decurrencies; leaves trifoliolate, sparsely pilose sometimes with hairs concentrated on midribs; petioles 1–1.5 mm long, glandular verrucose; terminal leaflets narrowly elliptic to oblanceolate, 2.5–5 × 1–3 mm, margins slightly to obviously glandular dentate, adaxial surface not glandular verrucose; inflorescence usually longer than the subtending leaves, 1–3-flowered; peduncle 3–5 mm long, sparsely stellate hairy; pedicels 1–1.5 mm long, with 1 mm, abaxial surfaces glabrous or with a sparse indumentum of minute stellate hairs; petals c. 2.5 × 1.2 mm, adaxial surface sparsely pilose, abaxial surface with a dense stellate indumentum; gynoecium glabrous; cocci and seed not seen (Duretto and Forster 2008; PlantNET 2016).
2. *Zieria odorifera* subsp. *warrabahensis* is endemic to New South Wales (NSW) and is known only from a limited area in and near Warrabah National Park west of Armidale in northern NSW. It is found in both heath and *Eucalyptus prava/Callitris endlicheri* layered woodland on granite outcrops (Duretto and Forster 2008; PlantNET 2016; L. Copeland *in litt.* May 2016, November 2016, January 2017).
3. *Zieria odorifera* subsp. *warrabahensis* is known from only two herbarium collections made c. 2.7 km apart. Two surveys for this taxon in October 2006 determined that the subspecies is confined to a small area in Warrabah National Park and adjacent private property, and it is apparently absent from other areas of the park that contain apparently suitable habitat. The population contains c. 100 mature individuals (L. Copeland *in litt.* May 2016, November 2016, January 2017); the number of mature individuals is therefore estimated to be very low.
4. The distribution of *Zieria odorifera* subsp. *warrabahensis* is very highly restricted. The area of occupancy (AOO) and the extent of occurrence (EOO) are estimated to be 4–8 km². The AOO is based on 2 x 2 km grid cells, the scale recommended for assessing AOO by IUCN (2016). The EOO is reported as equal to AOO, despite the range of the species, measured by a minimum convex polygon containing all the known sites of occurrence, being less than AOO. This is to ensure consistency with the definition of AOO as an area within EOO, following IUCN Guidelines (2017).
5. Threats to *Zieria odorifera* subsp. *warrabahensis* include grazing by feral goats and inappropriate fire regimes (L. Copeland, *in litt.* May 2016, January 2017). During

Established under the Biodiversity Conservation Act 2016

PO Box 1967 Hurstville BC NSW 1481

02 9585 6940 - Fax: 9585 6606

scientific.committee@environment.nsw.gov.au

NSW Threatened Species Scientific Committee

two separate surveys of the taxon across its entire range in October 2006, it was observed that a number of plants, both within and outside Warrabah National Park, had been heavily browsed with most leaves gone and just semi-broken stems remaining, and, as goats are prolific in the area, it was assumed they were responsible for this damage (L. Copeland, *in litt.* January 2017). The fire response of *Z. odorifera* subsp. *warrabahensis* has not been recorded but it is likely to resprout following fire given that this behaviour has been documented for New England populations of *Z. odorifera* (Clarke et al. 2009) (which can be assumed to be *Z. odorifera* subsp. *williamsii* based on geography). While individuals of *Z. odorifera* subsp. *warrabahensis* are therefore likely to survive a fire, repeated burns within a short interval may kill the plants, especially if their resources are already depleted as a result of frequent grazing (L. Copeland, *in litt.* January 2017). An absence of fire may limit population replacement unless the species can recruit new individuals in the absence of fire. The population's limited extent makes it highly susceptible to localised events such as clearing, disturbance and weed invasion. 'Competition and habitat degradation by Feral Goats, *Capra hircus* Linnaeus 1758', 'Clearing of native vegetation' and 'High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition' are listed as Key Threatening Processes under the Act.

6. *Zieria odorifera* subsp. *warrabahensis* Duretto & P.I.Forst. is eligible to be listed as a Critically endangered species as, in the opinion of the NSW Threatened Species Scientific Committee, it is facing a very high risk of extinction in Australia in the immediate future as determined in accordance with the following criteria as prescribed by the *Biodiversity Conservation Regulation 2017*:

Overall Assessment

Critically endangered under Clause 4.3

Clause 4.2 – Reduction in population size of species
(Equivalent to IUCN criterion A)

Assessment Outcome: Data Deficient

(1) – The species has undergone or is likely to undergo within a time frame appropriate to the life cycle and habitat characteristics of the taxon:			
	(a)	for critically endangered species	a very large reduction in population size, or
	(b)	for endangered species	a large reduction in population size, or
	(c)	for vulnerable species	a moderate reduction in population size.
(2) – The determination of that criteria is to be based on any of the following:			
	(a)	direct observation,	
	(b)	an index of abundance appropriate to the taxon,	
	(c)	a decline in the geographic distribution or habitat quality,	
	(d)	the actual or potential levels of exploitation of the species,	
	(e)	the effects of introduced taxa, hybridisation, pathogens, pollutants, competitors or parasites.	

NSW Threatened Species Scientific Committee

Clause 4.3 - Restricted geographic distribution of species and other conditions
(equivalent to IUCN criterion B)

Assessment Outcome: Critically endangered under Clause 4.3 (a) (d) (e i & iii).

The geographic distribution of the species is:			
	(a)	for critically endangered species	very highly restricted, or
	(b)	for endangered species	highly restricted, or
	(c)	for vulnerable species	moderately restricted,
and at least 2 of the following 3 conditions apply:			
	(d)	the population or habitat of the species is severely fragmented or nearly all the mature individuals of the species occur within a small number of locations,	
	(e)	there is a projected or continuing decline in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	habitat area, extent or quality,
		(iv)	the number of locations in which the species occurs or of populations of the species,
	(f)	extreme fluctuations occur in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	the number of locations in which the species occur or of populations of the species.

Clause 4.4 - Low numbers of mature individuals of species and other conditions
(Equivalent to IUCN criterion Clause C)

Assessment Outcome: Data Deficient

The estimated total number of mature individuals of the species is:			
	(a)	for critically endangered species	very low, or
	(b)	for endangered species	low, or
	(c)	for vulnerable species	moderately low,
and either of the following 2 conditions apply:			
	(d)	a continuing decline in the number of mature individuals that is (according to an index of abundance appropriate to the species):	
		(i)	for critically endangered species very large, or
		(ii)	for endangered species large, or
		(iii)	for vulnerable species moderate,
	(e)	both of the following apply:	
		(i)	a continuing decline in the number of mature individuals (according to an index of abundance appropriate to the species), and
		(ii)	at least one of the following applies:
		(A)	the number of individuals in each population of the species is:

NSW Threatened Species Scientific Committee

			(I)	for critically endangered species	extremely low, or
			(II)	for endangered species	very low, or
			(III)	for vulnerable species	low,
		(B)	all or nearly all mature individuals of the species occur within one population,		
		(C)	extreme fluctuations occur in an index of abundance appropriate to the species.		

Clause 4.5 - Low total numbers of mature individuals of species
(equivalent to IUCN criterion D)

Assessment Outcome: Endangered under Clause 4.5.

The total number of mature individuals of the species is:			
	(a)	for critically endangered species	extremely low, or
	(b)	for endangered species	very low, or
	(c)	for vulnerable species	low.

Clause 4.6 - Quantitative analysis of extinction probability
(Equivalent to IUCN criterion E)

Assessment Outcome: Data Deficient

The probability of extinction of the species is estimated to be:			
	(a)	for critically endangered species	extremely high, or
	(b)	for endangered species	very high, or
	(c)	for vulnerable species	high.

Clause 4.7 - Very highly restricted geographic distribution of species–vulnerable species

(Equivalent to IUCN criterion D2)

Assessment Outcome: Vulnerable via Clause 4.7

For vulnerable species,	the geographic distribution of the species or the number of locations of the species is very highly restricted such that the species is prone to the effects of human activities or stochastic events within a very short time period.
-------------------------	--

Dr Marco Duretto
Chairperson
NSW Threatened Species Scientific Committee

References:

Clarke PJ, Knox KJE, Campbell ML, Copeland LM (2009) Post-fire recovery of woody plants in the New England Tableland Bioregion. *Cunninghamia* **11**, 221–239.

NSW Threatened Species Scientific Committee

Duretto MF, Forster PI (2008) New subspecies for *Zieria odorifera* J.A.Armstr. (Rutaceae) from northern New South Wales. *Austrobaileya* **7**, 681–690.

IUCN Standards and Petitions Subcommittee (2016) Guidelines for Using the IUCN Red List Categories and Criteria. Version 12. Prepared by the Standards and Petitions Subcommittee.

<http://www.iucnredlist.org/documents/RedListGuidelines.pdf>

PlantNET (The NSW Plant Information Network System) Royal Botanic Gardens and Domain Trust, Sydney. <http://plantnet.rbgsyd.nsw.gov.au> (accessed 24 November 2016).