

NSW Threatened Species Scientific Committee

Notice of Preliminary Determination

The NSW Threatened Species Scientific Committee, established under the *Biodiversity Conservation Act 2016* (the Act), has made a Preliminary Determination to support a proposal to list the orchid *Prasophyllum wilkinsoniorum* D.L.Jones as a CRITICALLY ENDANGERED SPECIES in Part 1 of Schedule 1 of the Act. Listing of Critically Endangered species is provided for by Part 4 of the Act.

How to make a submission

The NSW TSSC welcomes public involvement in the assessment process and places preliminary determinations on public exhibition on the NSW TSSC pages on the NSW Department of Climate Change, Energy, the Environment and Water (NSW DCCEEW) website. This public exhibition provides an opportunity for the public to comment on this preliminary determination as well as provide any additional information that is relevant to the assessment.

Postal submissions regarding this Preliminary Determination may be sent to:
Secretariat
NSW Threatened Species Scientific Committee
Locked Bag 5022
Parramatta NSW 2124.

Email submissions in Microsoft Word or PDF formats to:
scientific.committee@environment.nsw.gov.au

Submissions close 3 July 2026

What happens next?

After considering any submissions received during the public exhibition period the NSW TSSC will make a Final Determination and a notice will be placed on the NSW DCCEEW website to announce the outcome of the assessment. If the Final Determination is to support a listing, then it will be added to the Schedules of the Act when the Final Determination is published on the legislation website. www.legislation.nsw.gov.au.

Privacy information

The information you provide in your submission may be used by the NSW TSSC in the assessment to determine the conservation status and listing or delisting of threatened or extinct species, threatened populations and threatened or collapsed ecological communities or to assess key threatening processes.

The NSW TSSC may be asked to share information on assessments with NSW Government agencies, the Commonwealth Government and other State and Territory governments to collaborate on national threatened species assessments using a common assessment method and to assist in the management of species and ecological communities.

NSW Threatened Species Scientific Committee

If your submission contains information relevant to the assessment it may be provided to state and territory government agencies and scientific committees as part of this collaboration.

If you wish your identity and personal information in your submission to be treated as confidential you must:

- *request your name be treated as confidential, and*
- *not include any of your personal information in the main text of the submission or attachments so that it can be easily removed.*

Professor Angela Moles, FRSN
Chairperson
NSW Threatened Species Scientific Committee

NSW Threatened Species Scientific Committee

Public exhibition period: 10/04/2026 - 3/07/2026 Preliminary Determination

The NSW Threatened Species Scientific Committee, established under the *Biodiversity Conservation Act 2016* (the Act), has made a Preliminary Determination to support a proposal to list the orchid *Prasophyllum wilkinsoniorum* D.L.Jones as a CRITICALLY ENDANGERED SPECIES in Part 1 of Schedule 1 of the Act. Listing of Critically Endangered species is provided for by Part 4 of the Act.

Summary of Conservation Assessment

Prasophyllum wilkinsoniorum D.L.Jones was found to be Critically Endangered in accordance with the following provisions in the *Biodiversity Conservation Regulation 2017*: Clause 4.3(a)(d)(e i,iii) because: (1) it has a very highly restricted geographic distribution with an estimated extent of occurrence of 12–24 km², (2) the minimum number of threat-defined locations when considering the most serious plausible threat of herbivory and habitat degradation by feral pigs is one (range 1–3), and (3) there is an observed and inferred continuing decline in habitat quality from anthropogenic habitat degradation and widespread feral pig diggings, as well as damage to potential habitat by domestic cattle, and an inferred continuing decline in the number of mature individuals from damage or consumption by pigs, deer, and rabbits, as well as anthropogenic habitat degradation.

The NSW Threatened Species Scientific Committee has found that:

1. *Prasophyllum wilkinsoniorum* D.L.Jones (Wilkinson's leek orchid, family Orchidaceae, CHAH 2018; PlantNET 2025) was described in Jones (2021) as: "Plants 250–450 mm tall. Free part of leaf to 200 x 4–8 mm. Spike 80–150 mm long, 15–45-flowered. Ovary shortly stalked, brownish. Flowers moderately crowded, scented, 15–19 x 12–15 mm, dark greenish-brown to dark brownish-red, labellum greenish to greenish-pink, callus green to brown. Dorsal sepal 8–11 x 3.5–4.5 mm, decurved. Lateral sepals free, 8–11 x 2 mm, widely spreading, margins pale. Labellum stalked, broadly ovate-lanceolate, 7.5–8.5 x 4.5–5.5 mm, sharply recurved near middle, constricted above middle then tapered to apex. Callus broad, with raised margins, extending nearly to labellum apex".
2. There are 2–4 known subpopulations of *Prasophyllum wilkinsoniorum*. These occur in the Kybeyan-Gourock subregion of the South Eastern Highlands (IBRA region, Commonwealth DCCEEW 2024a, 2024b), north-east of Bombala in New South Wales (NSW). The minimum number excludes the Hains Swamp subpopulation, which may no longer be extant, and the Tantawangalo subpopulation, which could not be distinguished from *P. canaliculatum* at the time of survey due to a lack of diagnostic reproductive material.
3. *Prasophyllum wilkinsoniorum* has an estimated total population size of approximately 2,080 mature individuals (see Smith 2025 for details and assumptions). In January 2025, ~200 mature individuals were detected in a survey that did not cover all potential habitat, nor the entire flowering period of this species.

NSW Threatened Species Scientific Committee

The observations of population dynamics for *P. correctum* (Coates *et al.* 2006) were applied to derive a preliminary estimate of population size for *P. wilkinsoniorum*. A precautionary approach to this estimation was applied given that the actual flowering rate for this species may differ from *P. correctum*, and there is an observed continuing decline in habitat quality alongside an inferred decline in the number of mature individuals for *P. wilkinsoniorum*.

4. *Prasophyllum wilkinsoniorum* has an estimated extent of occurrence (EOO) of 12–24 km², and an estimated area of occupancy (AOO) of 12–20 km². As recommended by IUCN (2024), AOO is based on 2 x 2 km grid cells, while EOO is based on a minimum convex polygon enclosing a cleaned dataset of current records for the species (ALA 2024; C. Steele *in litt.* December 2024; NSW Government 2024; RBGDT 2024; K.J. Smith, T.L. Collins and J. Miles pers. obs. January 2025; A. Morrison *in litt.* March 2025). The minimum EOO was adjusted from 4.6 km² to match the minimum AOO because, by definition, EOO cannot be less than AOO (IUCN 2024).
5. Jones (2021) described *Prasophyllum wilkinsoniorum* as “restricted to montane areas growing among tussocks and low shrubs in snowgum woodland, grassy flats, treeless plains, frost hollows and among sedges and rushes in swamps in brown basaltic loam, dark loam and peaty loam”. The species flowers from late December to January (Jones 2021). Individuals that emerge in the vegetative (*i.e.*, leaf only) state appear grass-like, and as such, may be easily missed during surveys (Coates *et al.* 2006). Some individuals may not emerge at all, instead remaining dormant underground (NSW NPWS 2022), possibly due to waterlogging, drought, or competition from dense growth of grasses and sedges (Miles 2017, Miles 2019, Miles 2024). From observations of *P. correctum*, it is inferred that *P. wilkinsoniorum* individuals can remain dormant for 1–5 years, and some mortality occurs among dormant tubers (Coates *et al.* 2006).
6. Based on the life history of other orchid species (Coates *et al.* 2006; Shefferson *et al.* 2020), and the small population size of *Prasophyllum wilkinsoniorum*, it is inferred that very few *P. wilkinsoniorum* seeds become mature adults. Assuming a seed is deposited in an area of suitable vegetation, soil, and climate, like other orchids, *Prasophyllum* species also require the presence of specific types of mycorrhizal fungi for germination and growth (Grant and Koch 2003; Freestone *et al.* 2023).
7. The lifespan of *Prasophyllum* species is uncertain; some deaths occur due to outside factors (*e.g.*, unsuitable environment, herbivory) but intrinsic mortality is poorly understood (Benson and McDougall 2005). For *P. wilkinsoniorum* it is inferred that a single dropper is produced each year to form a replacement tuber (Pridgeon and Chase 1995), which is considered the same individual. The generation length of *P. wilkinsoniorum* is uncertain; estimates of generation length for other *Prasophyllum* species range between 10–50 years (Commonwealth TSSC 2014; DELWP 2021, DELWP 2022; Commonwealth of Australia 2023).

NSW Threatened Species Scientific Committee

8. The major threats to the persistence of *Prasophyllum wilkinsoniorum* are herbivory and habitat degradation by feral animals and domestic cattle, anthropogenic habitat degradation, and competition with exotic plants. The resilience of *P. wilkinsoniorum* to these threats may be reduced by human-induced changes to rainfall and fire regimes. 'Predation, habitat degradation, competition and disease transmission by feral pigs, *Sus scrofa* Linnaeus 1758', 'Herbivory and environmental degradation caused by feral deer', 'Competition and grazing by the feral European Rabbit, *Oryctolagus cuniculus* (L.)', 'Clearing of native vegetation', 'Invasion of native plant communities by exotic perennial grasses', 'Loss and degradation of native plant and animal habitat by invasion of escaped garden plants, including aquatic plants', 'Anthropogenic climate change', and 'High frequency fire resulting in the disruption of life cycle processes in plants and animals and loss of vegetation structure and composition' are listed as Key Threatening Processes under the Act.
9. Continuing decline in habitat quality from extensive feral pig diggings was observed in the Nunnock, Dragon Swamp and Tantawangalo subpopulations of *Prasophyllum wilkinsoniorum* (K.J. Smith pers. obs. January 2025). There is also an inferred continuing decline in the number of mature individuals of *P. wilkinsoniorum* based on damage to, or consumption of, stems and tubers associated with feral animal activities. Targeting of *P. wilkinsoniorum* by feral pigs appears to have occurred in one of the Nunnock sites (Smith 2025), and adverse effects from the activities of deer and rabbits are suspected given their presence (McPherson 2004), and observations of rabbits digging up *Caladenia saggicola* (sagg spider orchid) tubers (Threatened Species Section 2017). Grazing and trampling of potential habitat by domestic cattle was evident in the paddocks adjoining the northern side of the road in the Hains Swamp locality where the persistence of *P. wilkinsoniorum* is currently unknown. Damage to flowering stems adversely affects an individual's ability to produce seed in a given year (McPherson 2004). Consecutive years of limited seed production, together with mortality caused by exposure or consumption when tubers are unearthed, places subpopulations at increased risk of extinction (McPherson 2004). In addition, bare patches created by feral and domestic animals may facilitate the encroachment of exotic plants that compete with *P. wilkinsoniorum* (McPherson 2004).
10. There is an inferred continuing decline in habitat and the number of mature *Prasophyllum wilkinsoniorum* based on the direct anthropogenic threats that were present at two localities in January 2025. In one Nunnock site, vehicle tracks were present through a patch of 27 individuals and there were no individuals in the area where vehicles had torn up some ground vegetation and compacted the soil. At the Hains Swamp locality, a main road was constructed through the swamp, and excavated soil is present as mounds along the roadsides. Part of the roadside at Hains Swamp was bulldozed in 2020 to serve as a firebreak (G. Phillips *in litt.* November 2024). This contrasts with the historical description of the Hains Swamp locality: Around 1999–2000 it was "in really good condition, quite wet and with a dense cover. There was also good grassy verge on both sides of the road that supported several species of orchid, and other monocots" (D.L. Jones in Miles 2020b).

NSW Threatened Species Scientific Committee

11. Exotic plants may degrade *Prasophyllum wilkinsoniorum* habitat by competing for space and resources. *Rubus fruticosus* sp. agg. around Dragon Swamp was noted as likely to require long-term control (J. Bentley *in litt.* October 2020). The 2019/20 wildfire at Dragon Swamp was followed by an increase in weed populations, principally *Cirsium vulgare* and *Holcus lanatus* (Miles 2020a). *Senecio madagascariensis* is also present (A. Morrison pers. comm. March 2025). In January 2025, *Holcus lanatus* was present in all subpopulations but particularly dense in some of the Nunnock Swamp sites. The Hains Swamp locality is most at risk of weed incursion due to the main road that runs through it. *Rubus fruticosus* sp. agg. is also present at Hains Swamp (K.J. Smith pers. obs. January 2025).
12. *Prasophyllum wilkinsoniorum* is considered to occupy 1–3 threat-defined locations based on the most serious plausible threat of herbivory and habitat degradation by feral pigs. A single location is considered most plausible given that the core range of feral pigs can be many square kilometres (Wilson *et al.* 2023), which could result in simultaneous adverse effects across all sites containing *P. wilkinsoniorum*.
13. *Prasophyllum wilkinsoniorum* D.L.Jones is eligible to be listed as a Critically Endangered species as, in the opinion of the NSW Threatened Species Scientific Committee, it is facing an extremely high risk of extinction in Australia in the immediate future as determined in accordance with the following criteria as prescribed by the *Biodiversity Conservation Regulation 2017*:

Assessment against *Biodiversity Conservation Regulation 2017* criteria

The Clauses used for assessment are listed below for reference.

Overall Assessment Outcome:

Critically Endangered under Clause 4.3(a)(d)(e i,iii)

Clause 4.2 – Reduction in population size of species (Equivalent to IUCN criterion A)

Assessment Outcome: Data Deficient

(1) - The species has undergone or is likely to undergo within a time frame appropriate to the life cycle and habitat characteristics of the taxon:			
	(a)	for critically endangered species	a very large reduction in population size, or
	(b)	for endangered species	a large reduction in population size, or
	(c)	for vulnerable species	a moderate reduction in population size.
(2) - The determination of that criteria is to be based on any of the following:			
	(a)	direct observation,	
	(b)	an index of abundance appropriate to the taxon,	
	(c)	a decline in the geographic distribution or habitat quality,	
	(d)	the actual or potential levels of exploitation of the species,	
	(e)	the effects of introduced taxa, hybridisation, pathogens, pollutants, competitors or parasites.	

NSW Threatened Species Scientific Committee

**Clause 4.3 - Restricted geographic distribution of species and other conditions
(Equivalent to IUCN criterion B)**

Assessment Outcome: Critically Endangered under Clause 4.3(a)(d)(e i,iii)

The geographic distribution of the species is:			
	(a)	for critically endangered species	very highly restricted, or
	(b)	for endangered species	highly restricted, or
	(c)	for vulnerable species	moderately restricted,
and at least 2 of the following 3 conditions apply:			
	(d)	the population or habitat of the species is severely fragmented or nearly all the mature individuals of the species occur within a small number of locations,	
	(e)	there is a projected or continuing decline in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	habitat area, extent or quality,
		(iv)	the number of locations in which the species occurs or of populations of the species,
	(f)	extreme fluctuations occur in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	the number of locations in which the species occur or of populations of the species.

Clause 4.4 - Low numbers of mature individuals of species and other conditions

(Equivalent to IUCN criterion C)

Assessment Outcome: Data Deficient

The estimated total number of mature individuals of the species is:				
	(a)	for critically endangered species	very low, or	
	(b)	for endangered species	low, or	
	(c)	for vulnerable species	moderately low,	
and either of the following 2 conditions apply:				
	(d)	a continuing decline in the number of mature individuals that is (according to an index of abundance appropriate to the species):		
		(i)	for critically endangered species	very large, or
		(ii)	for endangered species	large, or
		(iii)	for vulnerable species	moderate,
	(e)	both of the following apply:		
		(i)	a continuing decline in the number of mature individuals (according to an index of abundance appropriate to the species), and	
		(ii)	at least one of the following applies:	

NSW Threatened Species Scientific Committee

		(A)	the number of individuals in each population of the species is:
		(I)	for critically endangered species extremely low, or
		(II)	for endangered species very low, or
		(III)	for vulnerable species low,
		(B)	all or nearly all mature individuals of the species occur within one population,
		(C)	extreme fluctuations occur in an index of abundance appropriate to the species.

Clause 4.5 - Low total numbers of mature individuals of species

(Equivalent to IUCN criterion D)

Assessment Outcome: Not met

The total number of mature individuals of the species is:		
	(a)	for critically endangered species extremely low, or
	(b)	for endangered species very low, or
	(c)	for vulnerable species low.

Clause 4.6 - Quantitative analysis of extinction probability

(Equivalent to IUCN criterion E)

Assessment Outcome: Data Deficient

The probability of extinction of the species is estimated to be:		
	(a)	for critically endangered species extremely high, or
	(b)	for endangered species very high, or
	(c)	for vulnerable species high.

Clause 4.7 - Very highly restricted geographic distribution of species–vulnerable species

(Equivalent to IUCN criterion D2)

Assessment Outcome: Vulnerable under Clause 4.7

For vulnerable species,	the geographic distribution of the species or the number of locations of the species is very highly restricted such that the species is prone to the effects of human activities or stochastic events within a very short time period.
-------------------------	--

Professor Angela Moles, FRSN

Chairperson

NSW Threatened Species Scientific Committee

Supporting Documentation:

Smith KJ (2025) Conservation Assessment of *Prasophyllum wilkinsoniorum* D.L.Jones (Orchidaceae). NSW Department of Climate Change, Energy, the Environment and Water.

NSW Threatened Species Scientific Committee

References:

- ALA (2024) *Prasophyllum wilkinsoniorum* occurrence download [Online]. Available at <https://doi.org/10.26197/ala.5aa68536-dbb4-494c-9cd0-c4b5499324a7> (accessed 21 November 2024)
- Benson D, McDougall L (2005) Ecology of Sydney plant species. *Cunninghamia* **9**, 16–212.
- CHAH (2018) Australian Plant Census (APC) *Prasophyllum wilkinsoniorum* [Online]. Council of Heads of Australasian Herbaria. Available at <https://id.biodiversity.org.au/tree/51786579/51410001> (accessed 17 January 2025)
- Coates F, Lunt ID, Tremblay RL (2006) Effects of disturbance on population dynamics of the threatened orchid *Prasophyllum correctum* D.L. Jones and implications for grassland management in south-eastern Australia. *Biological Conservation* **129**, 59–69.
- Commonwealth DCCEEW (2024a) Interim Biogeographic Regionalisation for Australia (IBRA), Version 7 (Regions) [Online]. Commonwealth of Australia and Department of Climate Change, Energy, the Environment and Water. Available at <https://datasets.seed.nsw.gov.au/dataset/interim-biogeographic-regionalisation-for-australia-ibra-version-7-regions> (accessed 2 July 2024)
- Commonwealth DCCEEW (2024b) Interim Biogeographic Regionalisation for Australia (IBRA), Version 7 (Subregions) [Online]. Commonwealth of Australia and Department of Climate Change, Energy, the Environment and Water. Available at <https://datasets.seed.nsw.gov.au/dataset/interim-biogeographic-regionalisation-for-australia-ibra-version-7-subregions> (accessed 7 February 2024)
- Commonwealth of Australia (2023) Conservation advice for *Prasophyllum uvidulum* (summer leek-orchid). Commonwealth Department of Climate Change, Energy, the Environment and Water.
- Commonwealth TSSC (2014) Conservation advice *Prasophyllum laxum* lax leek-orchid. Commonwealth Department of Climate Change, Energy, the Environment and Water, Threatened Species Scientific Committee.
- DELWP (2021) Threatened Species Assessment *Prasophyllum diversiflorum* Gorae leek-orchid. Victoria State Government Department of Environment, Land, Water and Planning.
- DELWP (2022) Threatened Species Assessment *Prasophyllum litorale* coastal leek-orchid. Victoria State Government Department of Environment, Land, Water and Planning.

NSW Threatened Species Scientific Committee

- Freestone M, Linde C, Swarts N, Reiter N (2023) Asymbiotic germination of *Prasophyllum* (Orchidaceae) requires low mineral concentration. *Australian Journal of Botany* **71**, 67–78.
- Grant CD, Koch J (2003) Orchid species succession in rehabilitated bauxite mines in Western Australia. *Australian Journal of Botany* **51**, 453–457.
- IUCN (2024) Guidelines for Using the IUCN Red List Categories and Criteria Version 16. International Union for Conservation of Nature Standards and Petitions Subcommittee. Available at <https://www.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed 17 April 2024)
- Jones DL (2021) ‘A Complete Guide to Native Orchids of Australia, third edition.’ (Reed New Holland Publishers Pty Ltd: Sydney, Australia; Auckland, New Zealand)
- McPherson P (2004) Targeted flora survey report *Prasophyllum canaliculatum* Kybeyan and Tantawangalo December - January 2003-4. NGH Environmental.
- Miles J (2017) Searches and monitoring of *Prasophyllum canaliculatum* 2017. Jackie Miles, Consultant Botanist, unpublished report to NSW DCCEEW.
- Miles J (2019) Searches and monitoring of *Prasophyllum canaliculatum* 2019. Jackie Miles, Consultant Botanist, unpublished report to NSW DCCEEW.
- Miles J (2020a) Searches and monitoring of *Prasophyllum canaliculatum* 2020–21. Jackie Miles, Consultant Botanist, unpublished report to NSW DCCEEW.
- Miles J (2020b) Nomination for listing species as critically endangered, endangered or vulnerable under the NSW Biodiversity Conservation Act, 2016 – *Prasophyllum caricetum*. NSW Threatened Species Scientific Committee.
- Miles J (2024) Searches and monitoring of *Prasophyllum canaliculatum* 2023. Jackie Miles, Consultant Botanist, unpublished report to NSW DCCEEW.
- NSW NPWS (2022) ‘Conservation Action Plan Summer leek orchid (*Prasophyllum canaliculatum*).’ (Environment and Heritage, Department of Planning and Environment: Parramatta)
- NSW Government (2024) NSW BioNet [Online]. Available at <https://www.environment.nsw.gov.au/topics/animals-and-plants/biodiversity/nsw-bionet> (accessed 21 November 2024)
- PlantNET (2025) The Plant Information Network System of The Royal Botanic Gardens and Domain Trust Version 2.0. Available at <https://plantnet.rbgsyd.nsw.gov.au/> (accessed 17 April 2025)

NSW Threatened Species Scientific Committee

Pridgeon AM, Chase MW (1995) Subterranean axes in tribe Diurideae (Orchidaceae): Morphology, anatomy, and systematic significance. *American Journal of Botany* **82**, 1473–1495.

RBGDT (2024) *Prasophyllum wilkinsoniorum* specimen records [dataset accessed 10 December 2024]. In 'NSW Herbarium specimen catalogue'. (Royal Botanic Gardens and Domain Trust: EMu [RBGNSW] Application).

Shefferson RP, Jacquemyn H, Kull T, Hutchings MJ (2020) The demography of terrestrial orchids: life history, population dynamics and conservation. *Botanical Journal of the Linnean Society* **192**, 315–332.

Threatened Species Section (2017) Listing statement for *Caladenia saggicola* (sagg spider-orchid). Department of Primary Industries, Parks, Water and Environment.

Wilson C, Gentle M, Marshall D (2023) Factors influencing the activity ranges of feral pigs (*Sus scrofa*) across four sites in eastern Australia. *Wildlife Research* **50**, 876–889.

Expert Communications:

Andrew Morrison, Project Officer Threatened Species, NSW National Parks and Wildlife Service.

Christina Steele.

David L. Jones, Botanist

Gavin Phillips, Team Leader Threatened Species Assessment, NSW Department of Climate Change, Energy, the Environment and Water (DCCEEW).

Jackie Miles, Consultant Botanist.

Joss Bentley, Senior Project Officer, NSW DCCEEW.

Tim Collins, Senior Project Officer Threatened Species Assessment, NSW DCCEEW.